arXiv:1412.1458v2 [math.NT] 18 May 2016

A REMARK ON CHEVALLEY'S AMBIGUOUS CLASS NUMBER FORMULAS

CHIA-FU YU

Abstract. In this note we remark that Chevalley's ambiguous class number formula is an immediate consequence of the Hasse norm theorem, the local and global norm index theorems for cyclic extensions.

1. Introduction

The ambiguous ideal classes are the Galois invariants of the class group of a cyclic extension of a number field. In [1] Chevalley gave a formula for the ambiguous class number. Besides Chevalley's original paper, one can also find for Chevalley's ambiguous class formula in Gras' book (see [3, p. 178, p. 180]). Lemmermeyer [4] supplied a modern and elementary proof as its proof does not seem to appear in most textbooks of algebraic number theory. In this note we observe that this formula can be also deduced very easily from the Hasse norm theorem, the local and global norm index theorems for cyclic extensions. The proofs of these standard theorems can be found in most textbooks of algebraic number theory, for example in Lang's book [2]. We should note that our proof is not completely independent of the proofs appeared before as the Q-machine of Herbrand is also applied in the proof of the local norm index theorem.

We now state the ambiguous class number formula. Consider a cyclic extension K/k of number fields with cyclic Galois group $G = \mathrm{Gal}(K/k) = \langle\sigma\rangle$ of generator σ. Denote by $\mathfrak{o}$ and $\mathfrak{O}$ the ring of integers of k and K, respectively. Let ∞ and ∞_r (resp. $\widetilde{\infty}$ and $\widetilde{\infty}_r$) denote the set of infinite and real places of k (resp. of K), respectively, and $\mathbb{A}_k$ (resp. $\mathbb{A}_K$) the adele ring of k (resp. K). Let $r_k : \widetilde{\infty} \to \infty$ denote the restriction to k. A real cycle is a cycle which is supported in the set of real places [2, p. 123]. We may identify a real cycle with its support, which is a subset of real places.

Suppose that $\widetilde{\mathfrak{c}}$ is a real cycle on K which is stable under the G-action. Denote by

$$\mathrm{Cl}(K,\widetilde{\mathfrak{c}}) := \frac{\mathbb{A}_K^\times}{K^\times \widehat{\mathfrak{O}}^\times K_\infty(\widetilde{\mathfrak{c}})^\times} \tag{1.1}$$

the narrow ideal class group of K with respect to $\widetilde{\mathfrak{c}}$, where $\widehat{\mathfrak{O}}$ is the profinite completion of $\mathfrak{O}$, and $K_\infty(\widetilde{\mathfrak{c}})^\times = \{a = (a_w) \in K_\infty^\times \mid a_w > 0 \quad \forall\, w|\widetilde{\mathfrak{c}}\}$. Similarly one defines $\mathrm{Cl}(k,\mathfrak{c})$ for any real cycle $\mathfrak{c}$ on k. The group G acts on the finite abelian group $\mathrm{Cl}(K,\widetilde{\mathfrak{c}})$. Its G-invariant subgroup $\mathrm{Cl}(K,\widetilde{\mathfrak{c}})^G$ is called the *ambiguous ideal class group* (with respect to $\widetilde{\mathfrak{c}}$).

Date: May 19, 2016.

Let $\mathfrak{c}$ be the real cycle on k such that $\infty_r - \mathfrak{c} = r_k(\widetilde{\infty}_r - \widetilde{\mathfrak{c}})$, and let $\mathfrak{c}_0 := r_k(\widetilde{\mathfrak{c}})$. One has $\mathfrak{c} = \mathfrak{c}_0 \infty_r^c$, where ∞_r^c is the set of real places of k which does not split completely in K. Let $N_{K/k}$ denote the norm map from K to k. The cycle $\mathfrak{c}$ is determined by the property $N_{K/k}(K_\infty(\widetilde{\mathfrak{c}})^\times) = k_\infty(\mathfrak{c})^\times$. Put $\mathfrak{o}(\mathfrak{c})^\times := \mathfrak{o}^\times \cap i_\infty^{-1}(k_\infty(\mathfrak{c})^\times)$, where $i_\infty : k^\times \to k_\infty^\times$ is the diagonal embedding. Denote by V_f the set of finite places of k. Let $e(v)$ denote the ramification index of any place w over $v \in V_f$.

Theorem 1.1. *One has*

$$\#\operatorname{Cl}(K,\widetilde{\mathfrak{c}})^G = \frac{\#\operatorname{Cl}(k,\mathfrak{c})\prod_{v\in V_f} e(v)}{[K:k][\mathfrak{o}(\mathfrak{c})^\times : \mathfrak{o}(\mathfrak{c})^\times \cap N_{K/k}(K^\times)]}. \tag{1.2}$$

When $\widetilde{\mathfrak{c}} = \widetilde{\infty}_r$, we get the restricted version of the formula stated in [3, p. 178]. When $\widetilde{\mathfrak{c}} = \emptyset$, using an elementary fact

$$\#\operatorname{Cl}(k,\infty_r^c) = \frac{h(k)\cdot 2^{|\infty_r^c|}}{[\mathfrak{o}^\times : \mathfrak{o}(\infty_r^c)^\times]},$$

we get the ordinary version of the formula stated in [3, p. 180].

2. Proof of Theorem 1.1

Define the norm ideal class group $N(K,\widetilde{\mathfrak{c}})$ by

$$N(K,\widetilde{\mathfrak{c}}) := \frac{N_{K/k}(\mathbb{A}_K^\times)}{N_{K/k}(K^\times \widehat{\mathfrak{O}}^\times K_\infty(\widetilde{\mathfrak{c}})^\times)}. \tag{2.1}$$

Consider the commutative diagram of two short exact sequences (by Hilbert's Theorem 90)

$$\begin{array}{ccccccccc} 1 & \longrightarrow & \mathbb{A}_K^{\times 1-\sigma}\cap U & \longrightarrow & U & \xrightarrow{N_{K/k}} & N_{K/k}(U) & \longrightarrow & 1 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 1 & \longrightarrow & \mathbb{A}_K^{\times 1-\sigma} & \longrightarrow & \mathbb{A}_K^\times & \xrightarrow{N_{K/k}} & N_{K/k}(\mathbb{A}_K^\times) & \longrightarrow & 1, \end{array} \tag{2.2}$$

where $U = K^\times \widehat{\mathfrak{O}}^\times K_\infty(\widetilde{\mathfrak{c}})^\times$. The snake lemma gives the short exact sequence

$$1 \longrightarrow \operatorname{Cl}(K,\widetilde{\mathfrak{c}})^{1-\sigma} \longrightarrow \operatorname{Cl}(K,\widetilde{\mathfrak{c}}) \longrightarrow N(K,\widetilde{\mathfrak{c}}) \longrightarrow 1 \tag{2.3}$$

as one has an isomorphism $\mathbb{A}_K^{\times 1-\sigma}/(\mathbb{A}_K^{\times 1-\sigma}\cap U) \simeq \operatorname{Cl}(K,\widetilde{\mathfrak{c}})^{1-\sigma}$. On the other hand we have the short exact sequence

$$1 \longrightarrow \operatorname{Cl}(K,\widetilde{\mathfrak{c}})^G \longrightarrow \operatorname{Cl}(K,\widetilde{\mathfrak{c}}) \longrightarrow \operatorname{Cl}(K,\widetilde{\mathfrak{c}})^{1-\sigma} \longrightarrow 1, \tag{2.4}$$

which with (2.3) shows the following result.

Lemma 2.1. *We have* $\#\operatorname{Cl}(K,\widetilde{\mathfrak{c}})^G = \#N(K,\widetilde{\mathfrak{c}})$.

Define

$$\operatorname{Cl}(k,\mathfrak{c},\mathfrak{O}) := \frac{\mathbb{A}_k^\times}{k^\times k_\infty(\mathfrak{c})^\times N_{K/k}(\widehat{\mathfrak{O}}^\times)}.$$

Lemma 2.2. *The group* $N(K,\widetilde{\mathfrak{c}})$ *is isomorphic to a subgroup* $H \subset \operatorname{Cl}(k,\mathfrak{c},\mathfrak{O})$ *of index* $[K:k]$.

PROOF. Put $A := N_{K/k}(\mathbb{A}_K^\times)$, $B := N_{K/k}(K^\times \widehat{\mathfrak{O}}^\times K_\infty(\widetilde{\mathfrak{c}})^\times)$, $C := k^\times$ and $H := CA/CB$. The group H is a subgroup in $\mathrm{Cl}(k, \mathfrak{c}, \mathfrak{O})$, which is of index $[K : k]$ by the global norm index theorem [2, p. 193]. One has $A \cap C = N_{K/k}(K^\times) \subset B$ by the Hasse norm theorem [2, p. 195]. The lemma follows from

$$N(K, \widetilde{\mathfrak{c}}) = A/B = A/(A \cap C)B \simeq CA/CB = H. \quad \blacksquare$$

Consider the exact sequence

(2.5)

$$1 \longrightarrow \frac{\mathfrak{o}(\mathfrak{c})^\times}{\mathfrak{o}(\mathfrak{c})^\times \cap N(\widehat{\mathfrak{O}}^\times)} \longrightarrow \frac{\widehat{\mathfrak{o}}^\times}{N(\widehat{\mathfrak{O}}^\times)} \longrightarrow \mathrm{Cl}(k, \mathfrak{c}, \mathfrak{O}) \longrightarrow \mathrm{Cl}(k, \mathfrak{c}) \longrightarrow 1.$$

It is easy to see $\mathfrak{o}(\mathfrak{c})^\times \cap N_{K/k}(\widehat{\mathfrak{O}}^\times) = \mathfrak{o}(\mathfrak{c})^\times \cap N_{K/k}(K^\times)$ from the Hasse norm theorem. The local norm index theorem [2, p. 188, Lemma 4] gives

$$\# \left(\frac{\widehat{\mathfrak{o}}^\times}{N(\widehat{\mathfrak{O}}^\times)} \right) = \prod_{v \in V_f} e(v). \tag{2.6}$$

Combining Lemma 2.2, (2.5) and (2.6) we get

$$\#N(K, \widetilde{\mathfrak{c}}) = \frac{\#\,\mathrm{Cl}(k, \mathfrak{c}, \mathfrak{O})}{[K : k]} = \frac{\#\,\mathrm{Cl}(k, \mathfrak{c}) \prod_{v \in V_f} e(v)}{[K : k][\mathfrak{o}(\mathfrak{c})^\times : \mathfrak{o}(\mathfrak{c})^\times \cap N_{K/k}(K^\times)]}. \tag{2.7}$$

Theorem 1.1 follows from Lemma 2.1 and (2.7). $\blacksquare$

Remark 2.3. We do not know whether $\mathrm{Cl}(K, \widetilde{\mathfrak{c}})^G$ and $N(K, \widetilde{\mathfrak{c}})$ are isomorphic as abelian groups or whether there is a natural bijection between them. When $[K : k] = 2$ and $\#\,\mathrm{Cl}(K, \widetilde{\mathfrak{c}})^{1-\sigma}$ is odd, we show that there is a natural isomorphism

$$N(K, \widetilde{\mathfrak{c}}) \simeq \mathrm{Cl}(K, \widetilde{\mathfrak{c}})^G. \tag{2.8}$$

The map $1 - \sigma : \mathrm{Cl}(K, \widetilde{\mathfrak{c}}) \to \mathrm{Cl}(K, \widetilde{\mathfrak{c}})^{1-\sigma}$ restricted to $\mathrm{Cl}(K, \widetilde{\mathfrak{c}})^{1-\sigma}$ is the squared map Sq, which is an isomorphism from our assumption. The inverse of Sq defines a section of (2.4), and hence an isomorphism $\mathrm{Cl}(K, \widetilde{\mathfrak{c}}) \simeq \mathrm{Cl}(K, \widetilde{\mathfrak{c}})^G \oplus \mathrm{Cl}(K, \widetilde{\mathfrak{c}})^{1-\sigma}$. The assertion (2.8) then follows.

ACKNOWLEDGMENTS

The present work is done while the author's stay in the Max-Planck-Institut für Mathematik. He is grateful to the Institut for kind hospitality and excellent working environment. The author is partially supported by the grants MoST 100-2628-M-001-006-MY4 and 103-2918-I-001-009.

REFERENCES

[1] C. Chevalley, Sur la théorie du corps de classes dans les corps finis et les corps locaux, *J. Faculty Sci. Tokyo* 2 (1933), 365–476.

[2] S. Lang, *Algebraic number theory.* Graduate Texts in Mathematics, **110**. Springer-Verlag, New York, 1986. 354 pp.

[3] G. Gras, *Class field theory. From theory to practice.* Springer-Verlag, 2003. 491 pp.

[4] F. Lemmermeyer, The ambiguous class number formula revisited. *J. Ramanujan Math. Soc.* **28** (2013), no. 4, 415–421.

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA AND NCTS (TAIPEI OFFICE), ASTRONOMY MATHEMATICS BUILDING, NO. 1, ROOSEVELT RD. SEC. 4, TAIPEI, TAIWAN, 10617

E-mail address: chiafu@math.sinica.edu.tw